



Express Biotech International
4650 Wedgewood Blvd., STE 103
Frederick, MD 21703 USA

Tel: +001.301.228.2444
Fax: +001.301.560.6570
Email: xpressbio@xpressbio.com

Simian Reagents

Chagas (*Trypanosoma cruzi*): The antigen is a mixture of purified inactivated whole organism. It is a mixture of three strains from different geographical areas (US, Brazil, Argentina). It will cross react with Leishmania.

Chick: (Chikungunya) The antigen is a mixture of recombinant antigens: capsid, E1, and E2. It will cross react with other Alphavirus.

CMV (Cytomegalovirus): The antigen is a purified inactivated whole virus with group specific antigens. Most animals will have antibodies to this virus.

Dengue: The antigen is a mixture of purified inactivated whole virus. It is a mixture of four strains (Type 1, 2, 3, and 4). It may cross react with other Flavivirus.

Filovirus: The antigen is a mixture of recombinant antigens: glycoprotein from Zaire Eblola virus and Angola Marburgvirus.

SFV (Simian Foamy Virus): The antigen is a purified inactivated whole virus with group specific antigens. Most animals will have antibodies to this virus.

Hepatitis A: Due to the need for a sensitive assay, the assay protocol will be different for this assay. This HAV-IgG ELISA kit is based on solid phase, one-step incubation competitive principle ELISA method. If HAV IgG antibodies are present in the sample, they compete with monoclonal HAV-IgG antibodies labeled with horseradish peroxidase (HRP-Conjugate) for a fixed amount of purified native HAV antigens pre-coated in the wells.

Hepatitis B: Due to the need for a very sensitive and quantitative assay, the assay protocol will be different for this assay. For detection of anti-HBs, this kit uses antigen "sandwich" ELISA method where polystyrene microwell strips are pre-coated with recombinant HBsAg. Serum or plasma sample is added to the microwells together with a second HBsAg conjugated to Horseradish Peroxidase (HRP-Conjugate). In case of presence of anti-HBs in the sample, the pre-coated and conjugated antigens will be bound to the two variable domains of the antibody and during incubation, the specific immunocomplex formed is captured on the solid phase.

Herpes 1: The antigen is a recombinant gG HSV-1 protein. It is type specific for HSV-1 and will not cross react with HSV-2.

Herpes 2: The antigen is a recombinant gG HSV-2 protein. It is type specific for HSV-2 and will not cross react with HSV-1.

Herpes B Virus (Cercopithecine Herpesvirus 1): Due to the hazards associated with Herpes B Virus, the antigen HSV-1 (Herpes Simplex Virus Type 1) is used as a surrogate marker. It is a purified inactivated whole virus. It will cross react with HSV-2 (Herpes Simplex Virus Type 2) and other Herpes viruses.

Malaria: The antigen is a mixture of recombinant *Plasmodium falciparum* MSP1 protein and a recombinant *Plasmodium vivax* MSP1 protein. Please note that serology is not the gold standard for diagnosis of malaria. Other modalities along with serology should be used for suspected infected animals.

Measles: The antigen is a purified inactivated whole virus.

Mumps: The antigen is a purified inactivated whole virus.

Rabies: The antigen is a recombinant 34 kd phosphoprotein.

Rubella: The antigen is a purified inactivated whole virus

SARS-CoV-2 Nucleocapsid Protein (Wuhan Hu-1): A recombinant nucleocapsid protein (strain Wuhan Hu-1) produced in E. coli, encompassing amino acids 1-419.

SARS-CoV-2 RBD + M Protein (Wuhan Hu-1): A recombinant spike membrane fusion protein produces in HEK293 cells, which includes the receptor binding domain (RBD) of the SARS-CoV-2 spike protein (S-protein) and the virion surface domain of the membrane protein (M-protein) with a C-terminal HIS-tag.

SARS-CoV-2 Spike Protein (Omicron): The antigen is a full length trimeric spike protein (strain B1.1.529/Omicron) produced in HEK293 cells. Will cross react with other SARS-CoV-2 strains.

SARS-CoV-2 Spike Protein (Wuhan Hu-1): A mixture of recombinant SARS-CoV-2 Spike Glycoprotein (S1) and SARS-CoV-2 Spike Glycoprotein (S2) produced in HEK293 cells, sequence strain Wuhan Hu-1. Will cross react with other SARS-CoV-2 strains.

SIV (Simian Immunodeficiency Virus): The antigen is a purified inactivated whole virus. It will cross react with HIV (Human Immunodeficiency Virus).

St Louis Encephalitis Virus: The antigen is a recombinant NS-1 protein produced in 293 human cells. It will cross react with other Flavivirus.

SRV (Simian Retrovirus type D): The antigen is a purified inactivated whole virus with group specific antigens. Please note that the positive animals will normally have immunosuppression, thus the positive reactions will be weak on the assay. Positive animals will also show weak responses to other antigens.

STLV (Simian T Lymphotropic Virus): The antigen HTLV (Human T Lymphotropic Virus) is used as a surrogate marker. The antigen is a mixture of recombinant antigens: HTLV-1 gp21, HTLV-1 gp46, HTLV-2 gp 21 and HTLV-2 gp46.

SV40: (Simian Virus 40) The antigen is a purified inactivated whole virus. It will cross react with other polyoma viruses.



Express Biotech International
4650 Wedgewood Blvd., STE 103
Frederick, MD 21703 USA

Tel: +001.301.228.2444
Fax: +001.301.560.6570
Email: xpressbio@xpressbio.com

Toxoplasma: Toxoplasma gondii strain RH antigen is a purified inactivated whole organism propagated in mouse ascites.

VZV (Varicella Zoster Virus): The antigen is a purified inactivated whole virus with group specific antigens. Many animals will have antibodies to this virus.

West Nile Virus: The antigen is a mixture of purified inactivated whole virus. It is a mixture of two strains (New York and Uganda). It may cross react with other Flavivirus.

Yellow Fever: Yellow fever strain 17D: The antigen is a recombinant NS1 protein produced in 293 human cells.

Zika: The antigen is a recombinant NS-1 protein produced in 293 human cells. Sequence strain Uganda MR 766. It will cross react with other Flavivirus.

Primate ELISA reagents are for research use only, not for diagnostic use.